

**United States Department of Agriculture
Animal and Plant Health Inspection Service
Center for Veterinary Biologics
P. O. Box 844
Ames, IA 50010**

1. **Reagent Name:** *Leptospira canicola* Challenge Culture
2. **Strain or Source:** Moulton strain, NVSL No. 11203
3. **Lot Number:** Identified by date of preparation
4. **Fill Date:** Cultures are prepared every 6 to 16 weeks from *L. canicola*-infected hamsters.
5. **Expiration Date:** Use within 1 year from labeled date.

Precautions: *Leptospira* spp. are classified as a biosafety level 2 pathogen. Handle under biological safety hood while wearing gloves. Use face shield when inoculating animals.

6. **Intended Use:** Challenge material for *L. canicola* testing.
7. **Instructions for Use:** Upon receipt, subculture in fresh semisolid P80-BA medium before inoculating hamsters as described in Supplemental Assay Method (SAM) 609. *L. canicola* should be transferred through hamsters a minimum of three times before initiating a potency test according to SAM 609.
8. **Test of Reagent:** Cultures are obtained from infected hamsters challenged with *L. canicola* according to **SAM 609**. Testing results indicate that 10 to 100 organisms constitute 1 hamster LD₅₀. Cultures are negative for extraneous bacteria when incubated on blood agar plates at 37°C. Cultures are checked for serological identity.
9. **Container Size, Type, Weight, or Volume:** 16-mm x 125-mm screw-capped tubes containing 10 mL P80-BA semisolid medium with viable *L. canicola*.
10. **Storage Conditions:** Store at room temperature (20°- 25°C). After subculture, store at room temperature after dinger zones (foci of leptospiral growth) are apparent.
11. **CVB Technical Contact:** Bacteriology Section, Center for Veterinary Biologics, (515) 337-6140 or FAX (515) 337-7673.

12. Origin and Passage History: Origin: Commercial laboratory in 1969. This strain is hamster virulent.

13. Method of Preparation: Liver tissue from a *L. canicola*-infected hamster is homogenized and diluted in phosphate-buffered saline containing 1% bovine serum albumin. Dilutions of liver homogenate are inoculated onto semisolid P80-BA medium and incubated at 28°- 30°C for 2-4 weeks, or until dinger zones are apparent. Cultures are then stored at room temperature (20°- 25°C).

14. Other: None.

Reagent orders and feedback should be sent *including phone number* to the following email address: CVB@aphis.usda.gov

Reagent orders forms (APHIS 2018) are available from:
http://www.aphis.usda.gov/animalhealth/cvb_forms

REVISED: 15Apr14 alb